

Correlation Studies of three different F₂ Population in Rice (*Oryza sativa L.*)

SHAHINA PERWEEN¹, NISAR AKHTAR^{*2}, KRISHNA PRASAD¹, SURYA PRAKASH¹ AND EKHAQUE AHMAD¹

ABSTRACT

This study was performed to determine the association between seventeen agronomic traits of three F₂ population viz; Dhanrashi x Wita-12, Dhanrashi x Shyamala and Sahbhagidhan x Birsamati in rice at Birsa Agricultural University , Kanke, Ranchi, Jharkand in 2019. Sixteen characters were assessed in this study viz days to flowering, plant height, panicles per plant, panicle length, primary branches per panicle, secondary branches per panicle, spikelets per panicle, grains per panicle, 100 grain weight, total yield per plant, grain yield per plant, unfilled grains per plant, grains per plant, spikelets per plant, grain length and grain breadth. Results showed that all the characters except days to flowering and 100 grain weight were positively and significantly correlated with grain yield per plant and grains per panicle in all F₂ populations. grain yield per plant was negatively and less associated with grain length and L/B ratio in both Dhanrashi x Wita-12 and Sahbhagidhan x Birsamati where as in Dhanrashi x Shyamala grain yield is negatively associated with only L/B ratio.

Keywords: association ,rice, F₂ population, agronomic traits

INTRODUCTION

Rice [(*Oryza sativa L.*) (2n = 2x = 24)] is second most widely grown cereal crop. It is the staple food of more than half the world's population and grows in a wide range of environments. Worldwide, it is grown on an area of 166.1 million hectares with yield of 745.2 million tonnes (Anonymous, 2013). In India, rice covered an area of 44.2 million hectares during 2017 with production of 106.19 million tones and productivity of 2416 kg per hectare (Directorate of Economics & Statistics). The relationship between yield and its main economic components, in segregating populations of rice, has been studied by several researchers (Yogameenakshi and Vivekanandan, 2010). Selection of promising genotypes, in a breeding program, is based on various criteria, most importantly final crop yield and its quality. Yield is a complex character and composed of several components. Some of which affect the yield directly while others contribute indirectly. Correlation studies provide an opportunity to study the magnitude and direction of association of yield with its direct and indirect components and also among various components. Correlation analysis thus help in identifying suitable selection criteria for improving the yield. So, the present study was undertaken to know the inter-relation among different yield contributing characters and their association with grain yield. This study reveals the possibility of effective selection for improvement of yield and its important components in subsequent segregating populations.

MATERIALS AND METHODS

The present investigation was carried out during Kharif 2018 at rice experimental area of the Birsa Agricultural University, Ranchi, India. The experimental material comprised of five

parental lines (Dhanrashi, Wita-12, Shyamala, Sahbhagidhan and Birsamati) and F₂ population of three crosses involving these parents viz., Dhanrashi x Wita-12, Dhanrashi x Shyamala, Sahbhagidhan x Birsamati and the population size of these crosses were 309, 51 and 115 plants respectively. Each of the crosses represented a separate experiment. The F₂ materials were grown in the nursery on 25th June, 2018. Twenty two (22) days old seedlings were transplanted in the puddled field with a spacing of 25cm between rows and 20 cm between plants within a row. Recommended agronomic practices were followed throughout the crop growth period. Data were recorded on individual plants of each crosses for 17 important agronomic traits like days to flowering, plant height, panicles per plant, panicle length, primary branches per panicle, secondary branches per panicle, spikelets per panicle, grains per panicle, 100 grain weight, total yield per plant, grain yield per plant, unfilled grains per plant, grains per plant, spikelets per plant, grain length, grain breadth and L/B ratio .Statistical analyses, including estimation of descriptive statistics and coefficient of correlation was done for all three crosses.

RESULTS AND DISCUSSION

The success of plant breeding programs relies heavily on the existence of genetic variability in crops for a particular trait. Estimates for range, mean and coefficient of variation (CV) for all three F₂ populations are shown in Table. 1 . Among all agronomic traits, more phenotypic variations were found in panicles per plant, total yield per plant, grain yield per plant, unfilled grains per plant, grains per plant and spikelets per plant in all three (F₂) segregating populations, while days to flowering, panicle length, grain length and grain breadth had very less variations.

Different results was found in F₂ populations viz Dhanrashi x



ARTICLE INFO

Received on	:	09/08/2022
Accepted	:	16/09/2022
Published online	:	30/09/2022

Table1: Range, Mean and Coefficient of variation for different characters of three F₂ population of rice

Characters	Dhanrashi x Wita12			Dhanrashi x Shyamala			Sahbhagidhan x Birsamati		
	Range	Mean	CV%	Range	Mean	CV%	Range	Mean	CV%
Days to flowering	96-120	109.18	3.94	86-106	94.53	4.55	86-99	94.10	2.87
Plant height(cm)	50-187	106.95	14.07	66-128	102.22	13.42	70-136	109.12	11.52
Panicles per plant	1-19	8.92	40.69	3-21	11.67	37.78	2-20	9.86	41.27
Panicle length (cm)	17-32	27.44	8.03	21-31	26.75	7.94	21-32	27.23	7.71
Primary branches per panicle	8-19	14.51	13.86	7-17	12.59	14.49	8-17	12.13	14.59
Secondary branches per panicle	17-81	52.61	20.49	18-68	44.33	21.95	21-68	40.67	23.64
Spikelets per panicle	87-372	232.05	23.48	99-32	224.49	23.04	58-300	147.73	32.15
Grains per panicle	60-331	181.43	27.56	53-302	177.43	30.32	80-310	191.40	22.67
100 grain weight(g)	1.38-272	2.11	11.22	1.61-2.54	2.21	8.71	1.72-3.21	2.39	10.87
Total yield per plant(g)	3.62-69.66	26.67	49.77	5.73-64.63	32.43	47.08	3.03-67.09	24.53	51.15
Grain yield per plant(g)	2.65-62.08	23.89	52.12	5-60.13	29.61	48.80	2.86-60.15	22.10	52.16
unfilled grains per plant	15-950	320.49	51.93	75-590	300.16	45.38	15-745	244.49	65.46
Grains per plant	163.64-2664	1125.72	51.17	233.64-2811.94	1339.84	48.81	149.91-3105.31	1160.96	47.94
Spikelets per plant	222.92-3437.4	1446.22	46.39	308.64-3135.92	1640.00	45.94	134.91-2475.31	916.47	49.60
Grain length (mm)	5.5-9.5	7.24	9.16	6.5-7.7	7.16	3.87	7.1-11	8.41	10.12
Grain breadth(mm)	1.7-3.3	2.40	9.76	2.3-2.8	2.53	4.48	1.6-2.8	2.33	10.74
L/B ratio	2.07-4.56	3.06	15.73	2.32-3.35	2.84	6.95	2.63-5.75	3.67	18.22

Wita-12, Dhanrashi x Shyamala and Sahbhagidhan x Birsamati were presented in [table-2](#), [table-3](#) and [table -4](#) respectively. In Dhanrashi x Wita-12 days to flowering was highly correlated with only primary branches per panicle and grain length at both genotypic and phenotypic level. Where as plant height, panicle length, primary branches per panicle and secondary branches per panicle were positively and significantly associated with almost all the characters i.e spikelets per panicle, grains per panicle, total yield per plant ,grain yield per plant, unfilled grains per panicle, grains per plant and spikelets per plants except 100 grain weight, grain length, grain breadth and L/B ratio. ([Singh et al. 2018](#)) reported that the seed yield exhibited positive and significant correlation with plant height, effective tillers per plant and panicle length these characters are major yield contributing characters. Similar findings were also reported by ([Sravan et al. 2012](#), [Kalyan et al. 2017](#)). Spikelets per panicle and grains per panicle was highly correlated with almost all characters viz; total yield per plant ,grain yield per plant, unfilled grains per panicle, grains per plant and spikelets per plants except 100 grain weight and grain breadth.

Correlation was highly correlated between total yield per plant and grain yield per plant with almost all the characters except grain length, grain breadth and L/B ratio. ([Kharb et al. 2016](#)) reported positive correlation of grain yield with number of grains per panicle. Similar findings were also reported by ([Ramya Rathod et al. 2017](#), [Patel et al. 2018](#)). However in Dhanrashi x Shyamala results was found that positively significant correlation was observed between days to flowering with primary branches per panicle, secondary branches per panicle and spikelets per panicle. It was observed to be negatively correlated with the characters namely panicles per plant, 100 grain weight and grain breadth. Correlation coefficient of all the characters are non significant. Plant height, panicles per plant, panicle length, primary branches per panicle ,secondary branches per panicle, spikelets per panicle and grains per panicle were

highly correlated with almost all characters except 100 grain weight, grain length, grain breadth and L/B ratio similarly ([Ramya Rathod et al. 2017](#) and [Rachana Bagudam et al. 2018](#)) reported this type of results on their work. Panicles per plant was observed to be positively correlated with 100 grain weight, grain length and grain breadth but it was negatively correlated with L/B ratio having low magnitude of correlation coefficient at phenotypic level. In present study non significant correlation coefficient was observed between primary branches per panicle with 100 grain weight, grain length, grain breadth and L/B ratio. The grain yield per plant and total yield per plant was highly correlated with almost all characters except grain length, grain breadth and L/B ratio at phenotypic level which were supported by ([Kalyan et al. 2017](#)) whereas genotypically grain yield per plant was highly correlated with all characters except unfilled grains per plant, grains per plant and spikelets per plant. whereas in cross Sahbhagidhan x Birsamati days to flowering was significantly and positively correlated with grains per panicle, 100 grain weight, total yield per plant, grain yield per plant ,spikelets per plant and grain breadth at genotypic level but results are different at phenotypic level. Negative correlation for days to flowering with plant height, panicle per plant, unfilled grains per plant and grains per plant was observed. Plant height was observed to be positively and significantly associated with total yield per plant, grain yield per plant and spikelets per plant,panicle per plant ,panicle length, primary branches per panicle, secondary branches per panicle, unfilled grains per plant and grains per plant at phenotypic level , this type of findings was also observed by ([Rachana Bagudam et al. 2018](#), [Patel et al. 2018](#), [Ramya Rathod et al. 2017](#), [Kalyan et al. 2017](#), [Sameera et al. 2016](#), [Basavaraja et al. 2013](#), [Sravan et al. 2012](#)), Phenotypically primary branches per panicle, secondary branches per panicle, spikelets per panicle and grains per panicle was highly correlated with almost all characters except 100 grain weight, unfilled grains per plant, grain length.

Table 2: Genotypic and Phenotypic Correlation between different characters in Dhanrashi X Wita-12

Characters	G	P	Plant height	Panicles per Plant	Panicle length	Primary Branches per Panicle	Secondary Branches per Panicle	Spikelets per panicle	Grains per panicle	100 Grain Weight	Total yield per plant	Grain yield per plant	Unfilled grains per plant	Grain s Per Plant	Spikelets Per Plant	Grain Length h	Grain Breadth h	L/B ratio			
Days to flowering	G	0.019	-0.182**	0.949**	0.243**	0.261**	0.289**	0.242**	-0.711**	-0.168**	0.066	-0.096	0.205*	*	-0.346**	0.626**	-0.770**	0.246**			
Plant height	G	P	-0.166**	-0.083	0.053	0.119*	0.078	0.021	-0.047	-0.130*	-0.143**	-0.133*	-0.026	0.111*	-	-0.102	0.211**	-0.035	-0.106		
Panicles per plant	G	P	0.130* 0.278**	0.498** 0.426**	0.379** 0.313**	0.385** 0.340**	0.397** 0.310**	0.734** 0.202**	0.059	0.174** 0.305**	0.164** 0.297**	0.164** 0.178**	0.166*	0.111*	*	0.185** 0.299**	0.103 0.109	-0.097 -0.072	0.149** 0.129*		
Panicle length	G	P														0.950*	0.999**	0.035	-0.065	0.055	
Primary Branches per Panicle	G	P														0.791*	0.819**	-0.041	-0.032	-0.004	
Secondary Branches per Panicle	G	P														0.340*	0.415**	0.040	-	0.165**	
Spikelets per panicle	G	P														0.482**	0.482**	*	0.470**	0.105	
Total yield per plant	G	P														0.367**	0.277**	*	0.458*	0.137*	
Grain yield per plant	G	P														0.460**	0.450**	*	0.472**	-0.108	
Unfilled grains per plant	G	P														0.350**	0.355**	*	0.426*	0.176**	
Grain Length	G	P														0.410**	0.397**	*	0.413*	0.076	
Grain Breadth	G	P														0.434**	0.423**	*	0.431**	-0.095	
Grains per plant	G	P														0.676**	0.679**	*	0.448*	0.141*	
Grain length	G	P														0.749**	0.707**	*	0.471**	0.166**	
Grain breadth	G	P														0.676**	0.593**	*	0.461*	0.380**	
Grains per panicle	G	P														0.742**	0.745**	*	0.472*	0.228**	
Grains per plant	G	P														-0.104	0.742**	*	0.269*	0.204**	
100 Grain weight	G	P														-0.006	0.503**	*	0.277*	0.972**	
Total yield per plant	G	P															0.343**	0.369**	*	0.178**	0.219**
Grain yield per plant	G	P															0.2272**	0.274**	0.026	0.072	0.206**
Unfilled grains per plant	G	P															0.961**	0.376**	*	0.992*	0.437**
Grains per plant	G	P															0.992**	0.516**	*	0.968*	-0.351**
Grain length	G	P															0.990*	0.390**	*	0.959**	0.008
Grain breadth	G	P															0.455**	0.455**	*	0.974*	-0.027
Grain length	G	P															0.974*	0.949**	*	0.947*	-0.025
Grain breadth	G	P															0.476*	0.513**	*	0.594**	0.028
Grains per plant	G	P															0.473*	0.654**	*	0.451**	-0.017
Spikelets per plant	G	P															0.595**	0.595**	*	0.131**	0.078
Grain length	G	P															0.976**	0.976**	*	0.051	0.058
Grain breadth	G	P															0.119**	0.188**	*	0.188**	0.069

** indicates significant at 1% and * indicates significant at 5% level

Table 3: Genotypic and Phenotypic correlation between different characters in Dhanrashi x Shyamala

Characters	G	P	Panicles per plant	Branches per Panicle	Primary Branches per Panicle	Second ary Branch es Panicle	Spikel lets per panicle	Grains per panicle	100 Grain Weight	Total yield per plant	Grain yield per plant	Unfilled grains per plant	Grains per Plant	Spikelets per Plant	Grain Length	Grain Breadth	L/B ratio
Days to flowering	G 0.019 P 0.013	- 0.177 0.026	0.213 0.007	0.012 0.414**	-0.114 0.242*	0.122 0.266*	0.451** 0.250	0.190 0.136	0.195 0.131	0.031 0.047	0.139 0.192	0.115 0.175	0.401** 0.441**	0.006 0.013	0.336* 0.018	0.006 0.061	
Plant height	G 0.064 P 0.340	- 0.201 0.404	0.147 0.483**	0.241 0.512**	0.346* 0.451**	0.453** 0.075	0.079 0.465**	0.048 0.444**	0.440** 0.482**	0.119 0.441**	0.031 0.470**	0.355* 0.112	-0.358** 0.470**	0.013 0.081	0.909** 0.081	0.013 0.061	
Panicles per plant	G	-	0.399	-0.823	-0.163	-0.439**	-0.574**	-0.754**	-0.621**	-	-0.279*	-0.604**	-0.478**	0.577**	-0.284*	0.529**	
Panicle length	G	-	0.365	0.138	0.371**	0.384**	0.367**	0.159	0.902***	0.650**	0.892**	0.686**	0.899**	0.000	0.103	-0.070	
Primary Branches per Panicle	G	-	0.528	0.204	0.591**	0.544**	0.482**	0.238	0.894**	0.622***	0.896**	0.563**	0.563***	0.391***	0.788**	0.000	0.324*
Secondary Branches per Panicle	G	-	0.027	-0.550**	-0.908**	0.367**	-0.178	-0.265	0.159	-0.349*	-0.328*	-0.328*	-0.328*	-0.274*	0.497**	-0.21	0.444**
Spikeslets per panicle	G	-	0.827**	0.671**	0.645**	0.011	0.392**	0.382**	0.378**	0.392**	0.408**	0.076	-0.404**	0.021	0.025	-0.021	0.025
100 Grain Weight	G	-	0.600**	-0.994**	0.572**	-0.231	-0.351*	0.381**	-0.351*	-0.394**	-0.404**	-0.404**	-0.404**	-0.912**	0.512**	-0.013	0.790**
Total yield per plant	G	-	0.771**	0.706**	-0.036	0.578**	0.564**	0.508**	0.508**	0.588**	0.602**	0.186	-0.186	-0.186	-0.013	0.102	-
Grains per panicle	G	-	0.396**	-0.005	0.538***	-0.721**	-0.298*	-0.124	-0.320*	-0.171	-0.096	-0.218	-0.390**	-0.390**	-0.078	0.144	-
Unfilled grains per plant	G	-	0.007	0.488**	0.480**	0.349*	0.480**	0.349*	0.510**	0.505**	0.090	-0.104	-0.104	-0.104	0.101	-	-
Grains per plant	G	-	0.794**	-	0.527**	-0.047	-0.777**	-0.666**	-0.666**	0.096	-0.606**	-0.606**	-0.606**	-0.393**	-	-	-
Spikeslets per Plant	G	-	0.233	0.231	0.154	0.039	0.062	0.062	0.062	-0.113	0.005	-0.057	-0.057	-0.057	-	-	-
Grain Length	G	-	0.993**	0.750**	-0.077	0.328*	-0.009	-0.061	-0.061	-0.951**	0.037	0.037	-0.525**	-0.525**	0.001	0.673**	-
Grain Breadth	G	-	0.695**	0.695**	0.201	-0.058	-0.832**	-0.832**	-0.832**	0.053	0.052	-0.672**	-0.672**	-0.672**	-0.014	-0.014	-
Spikeslets per Plant	G	-	0.682**	0.682**	0.486**	0.113	-0.733**	-0.733**	-0.733**	0.086	0.054	-0.537**	-0.537**	-0.537**	0.001	0.652**	-
Grains per Plant	G	-	0.091**	-	0.991**	-0.047	-0.193	-0.193	-0.193	0.086	0.054	-0.509**	-0.509**	-0.509**	0.001	-0.509**	-
Grain Breadth	G	-	0.085	0.085	-0.591**	-0.591**	-0.041	-0.041	-0.041	0.041	0.041	-0.420**	-0.420**	-0.420**	0.011	0.011	-
Grain Length	G	-	0.806**	-0.408**	-0.581**	-0.581**	-0.036	-0.036	-0.036	-0.036	-0.036	-0.873**	-0.873**	-0.873**	0.060	0.060	-
Grain Breadth	G	-	-0.244	-0.244	-0.867**	-0.867**	-	-	-	-	-	-0.867**	-0.867**	-0.867**	-0.001	-0.001	-

** indicates significant at 1% and * indicates significant at 5% level

Table-4: Genotypic and Phenotypic correlation between characters in Sahbhaghidhan X Birksamati

Characters	G / P	Panicle height	Panicles per plant	Panicle length	Primary Branches per Panicle	Secondary Branches Per Panicle	Spikele ts Per panicle	Grain s per Panicl e	100 Grain Weight	Total yield per plant	Grain yield per plant	Unfill ed grains per plant	Grain ns per Plant	Spikel ets per plant	Grain Length	Grain Breadt h	L/B ratio
Days to flowering	G P	0.750** -0.201**	-0.765** -0.124	0.005 0.057	-0.464** 0.097	-0.137 0.030	0.432** -0.111	0.792* *0.030	0.788** -0.063	0.280** -0.070	0.386 0.042	0.504** -0.050	-0.012 0.006	0.589** -0.131	-0.146 0.092	0.459** 0.092	
Plant height	G P	0.847** 0.231**	0.094 0.196*	0.449 ***	-0.539** 0.257**	-0.276** 0.123	0.744* *0.059	0.932** 0.273**	0.822** 0.265**	0.962** 0.265*	0.067 0.271	0.300** 0.258**	-0.446** 0.038	0.459 -0.010	0.311** 0.040	-0.146 0.092	
Panicles per plant	G P	0.132 0.226	0.144 0.102	0.156	0.275** 0.095	-0.257** 0.272*	0.597** *0.095	0.803** 0.180	0.871** 0.742**	0.565** 0.476**	0.741 0.740**	0.773** 0.740**	-0.411 -0.004	0.200** -0.118	-0.146 0.095	0.327** 0.095	
Panicle length	G P				-0.272** 0.355	-0.337** 0.381**	-0.551** 0.253**	0.633* *0.34*	0.784** 0.147	0.265** 0.372**	0.473** 0.677**	0.558 0.378	0.996** 0.373**	-0.425** 0.076	0.212** 0.070	-0.146 0.005	0.772** 0.005
Primary Branches per panicle	G P				0.847** 0.883**	0.581** 0.678**	0.636* *0.636*	0.749* *0.636*	0.569** -0.078	0.451** 0.245**	0.498** 0.254**	0.027 0.080	0.454** 0.284**	-0.207** 0.157	0.058 -0.205*	-0.166 0.217*	-0.146 0.168
Secondary branches per panicle	G P				0.683** 0.736**	0.616* 0.762*	0.550** 0.034	0.593** 0.343**	0.646** 0.348**	0.095 0.156	0.503 0.336	0.629** 0.356**	-0.212** 0.133	0.098 -0.143	-0.191** 0.168	-0.146 0.168	
Spikelets per panicle	G P				0.765* 0.661*	0.805** 0.019	0.354** 0.296**	0.413** 0.320**	0.318** -0.075	0.104 0.246	0.259** 0.328**	-0.265 0.154	0.078 -0.200*	-0.198* 0.223*	-0.146 0.223*	-0.146 0.223*	
Grains per panicle	G P					-0.296** 0.035	0.183* 0.385**	0.894** 0.384**	0.838** 0.201*	0.570 0.387	0.990** 0.403**	0.764** -0.020	-0.232* -0.108	0.076 -0.221*	-0.146 0.221*	-0.146 0.221*	
100 Grain Weight	G P						0.101 0.327**	0.026 0.339**	0.774** 0.115	0.443 0.161	0.289** 0.156	0.833 -0.012	-0.847** 0.396**	0.567** -0.251*	-0.146 0.251*	-0.146 0.251*	
Total yield per plant	G P							0.939** 0.994**	0.841** 0.596**	0.307** 0.068	0.975** 0.048	0.234* -0.048	-0.089 0.097	0.202* -0.075	-0.146 0.075	-0.146 0.075	
Grain yield per plant	G P								0.816** 0.519**	0.992 0.947	0.998** 0.977**	0.332** -0.031	-0.168 0.077	0.308** -0.053	-0.146 0.053	-0.146 0.053	
Unfilled grains Per plant	G P									0.866 0.723	0.800** 0.533**	0.555** -0.117	0.527** 0.165	-0.168 0.077	0.308** -0.053	-0.146 0.053	-0.146 0.053
Grains per plant	G P										1.000** 0.970**	0.970** 0.050	-0.140 0.028	0.130 -0.029	-0.146 0.051	-0.146 0.051	-0.146 0.051
Spikelets per plant	G P												-0.067 -0.041	-0.058 0.003	-0.146 0.011	-0.146 0.011	
Grain Length	G P												0.904** -0.312**	0.898** -0.762**	-0.146 0.465**	-0.146 0.465**	
Grain Breadth	G P												-	0.840** -	-0.146 0.840**	-0.146 0.840**	

** indicates significant at 1% and * indicates significant at 5% level

CONCLUSION

In summary, the results of this study indicated that days to flowering, 100 grain weight and grains per panicle could be considered as critical criteria for yield improvement in segregating generations of rice.

REFERENCES

- Anonymous. 2013. Available at <http://faostat.fao.org> (accessed on August 23, 2014).
- Basavaraja T, Asif M, Mallikarjun SK and Gangaprasad S.2013.Correlation and path analysis of yield and yield attributes in local rice cultivars (*Oryza sativa* L.). *Asian Journal of Bio Science* 8(1):36-38.
- Kalyan B, Radha Krishna KV and Subba Rao LV.2017. Correlation Coefficient Analysis for Yield and its Components in Rice (*Oryza sativa* L.) Genotypes. *International Journal of Current Microbiology and Applied Sciences* 6(7): 2425-2430.
- Kharb A, Jain S and Jain R K. 2016. Phenotypic analysis, correlation studies and linkage mapping of QTL for traits promoting cultivation under dry direct seeded aerobic conditions for the development of water-efficient high yielding rice lines. *Cereal Research Communications* 44(4):658-668.
- Patel HR, Patel VP, Patel PB, Rathod AJ and Pampaniya AG.2018. Genetic variability, correlation and path analysis for grain yield and component traits in F₃ segregating population of rice (*Oryza sativa* L.) *International Journal of Chemical Studies* 6(2):2327-2331.
- Rachana Bagudam KB, Eswari, Badri J and Raghuveer Rao P. 2018.Correlation and Path Analysis for Yield and its Component Traits in NPT Core Set of Rice (*Oryza sativa* L.). *International Journal of Current Microbiology and Applied Sciences* 7(9):97-108.
- Ramya Rathod D, Sanjeeva Rao V, Babu R and Bharathi M. 2017. Correlation and Path Coefficient Analysis for Yield, Yield Attributing and Nutritional Traits in Rice (*Oryza sativa* L.). *International Journal of Current Microbiology and Applied Sciences* 6(11): 183-188.
- Sameera T, Srinivas AP, Rajesh V, Jayalakshmi and Nirmala PJ.2016. Variability and path co-efficient for yield and yield components in rice. *Bangladesh Journal of Agricultural Research* 41(2): 259-271.
- Singh A, Singh A P, Singh A, Maurya M K, Yadav V, Singh H and Avasthi S. 2018. Study of simple correlation coefficients for yield and its component traits in rice (*Oryza sativa* L.). *Journal of Entomology and Zoology Studies* 6(4):1774-1777.
- Sravan T, Rangare NR, Suresh BG and Ramesh KS.2012. Genetic variability and character association in rainfed upland rice (*Oryza sativa* L.). *Journal of Rice Research* 5(1-2): 24-28.
- Yogameenakshi P and Vivekanandan P.2010. Association analysis in F1 and F2 generations of rice under reproductive stage drought stress. *Electron. J Plant Breed* 1(4):890-898.

ACKNOWLEDGMENTS

Authors are thankful to the Head, Scientists, Professors and technical staff, Department of Genetics and Plant Breeding, Birsa Agricultural University, Kanke Ranchi, Jharkhand for their valuable assistance.

Citation:

Perween S, Akhtar N, Prasad K, Prakash S and Ahmad E.2022. Correlation studies of three different F₂ population in Rice (*Oryza sativa* L.). *Journal of AgriSearch* 9(3):212-217